

KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY

**WARNING**

**DONAGHYS**

# Premium Bloat AF

An aid in the prevention and treatment of frothy bloat in cattle suitable for use via in-line dispensing systems.

Contains 880 g/L of Ethylene Oxide / Propylene Oxide Block Co-polymers

**DONAGHYS**  
PERFORMANCE FIRST

# DONAGHYS PREMIUM BLOAT AF

An aid in the prevention and treatment of frothy bloat in cattle suitable for use via in-line dispensing systems.

## WARNING

### READ LABEL BEFORE USE

### HANDLING PRECAUTIONS

Wear protective gear including gloves, protective clothing and eye protection. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. For animal treatment only.

### DIRECTIONS FOR USE

**By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on label.**

Start treatment before the risk becomes severe using a low dose rate to accustom stock to the treatment.

For in line dosing start dosing 3 weeks before the onset of a bloat challenge at a rate of 4mL/ cow every 24 hours. One week before the expected bloat challenge, increase the dose to 7mL. During the bloat challenge a dose rate of 10mL per cow every 24 hours must be used. These rates are for undiluted product.

For manual trough treatment Premium Bloat AF is added as a diluted premix twice per day. The premix is made by adding 1 part of Premium Bloat AF to 4 parts of water. Refer to table below for dose rates and timing. Pre-treat all water troughs with 150mL of premix to every 100 litres of water before regular dosing.

	3 weeks before onset of bloat	1 week before onset of bloat	During bloat challenge
	Premium Bloat AF per cow every 24 hours		
Dosing System	4mL	7mL	10mL
	1:4 Diluted mixture per cow every 12 hours		
Manual Trough Treatment	8mL	10mL	16mL
Drenching	-	-	25-50mL

**DRENCHING:** Dose each animal with 25 to 50mL of 1:4 premix, depending on severity of bloat, every 12 hours.

**SEVERLY BLOATED COWS:** Drench the animal with 300mL of 1:9 (30mL Premium Bloat AF and 270mL warm water) from a long neck bottle.

**WARNING:** Traces of detergent-based bloat remedies may be toxic to calves especially pre-weaning. Wash mixing buckets and containers thoroughly before reusing.

**HAZARD STATEMENT:** Causes serious eye irritation.

**RESPONSE STATEMENT: If in eyes:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do so. Continue rinsing. If eye irritation persists, get medical attention.

### STORAGE

Store in original container, tightly closed, away from foodstuffs, at temperatures below 25°C and out of reach of children.

### DISPOSAL

Dispose of empty containers safely at the nearest Agrecovery point or in accordance with local regulations. Triple rinse containers, add rinsings to the solution. Avoid contamination of any water supply with chemical or empty container. Dispose of chemicals safely in accordance with local regulations. Avoid contamination of any water supply.

ZPLAB757 MAR25 PN-036874-1 VER2

**DONAGHYS**  
PERFORMANCE FIRST

REGISTERED TO DONAGHYS LTD  
16 SHEFFIELD CRESCENT, CHRISTCHURCH, NEW ZEALAND  
FREEPHONE: 0800 942 006, WWW.DONAGHYS.COM  
EPA APPROVAL CODE: HSR002386  
REGISTERED PURSUANT TO THE ACVM ACT 1997 NO. A011688  
SEE WWW.FOODSAFETY.GOV.TZ FOR REGISTRATION CONDITIONS

