

FOR ANIMAL TREATMENT ONLY. KEEP OUT OF REACH OF CHILDREN.

# DONAGHYS Cobalt

Dairy cattle mineral supplement



Cobalt is an essential component of Vitamin B12 and thus is a key contributor to energy levels in the dairy cow. Overcoming low Cobalt levels can assist with:

- General good thrift
- Improved milk production
- Improved appetite
- Improved fertility

8.5mg/mL

**DONAGHYS**  
PERFORMANCE FIRST



# Donaghys Cobalt

Help your cows achieve a higher level of performance with Cobalt, the dairy cow liquid mineral supplement!

## WARNING

**READ LABEL BEFORE USE**

## HANDLING PRECAUTIONS

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves, protective clothing, eye and face protection. Wash hands thoroughly after handling. Avoid breathing mist/vapours/spray. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Collect spillage.

## HAZARD STATEMENTS

May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure. Toxic to aquatic life with long lasting effects.

## DIRECTIONS

Only for use when there is a confirmed Cobalt deficiency.  
Ready to use liquid no dilution required.

## DONAGHYS COBALT DOSE RATE

**DO NOT EXCEED THE STATED DOSE RATE AND FREQUENCY**

### Through water supply

Can be added to drinking water via dispenser. Administer to cattle daily at the following rate: 15mL per 100 animals per day. Do not exceed dose rate of 2mL per animal per day.

### Oral drenching

Administer orally to cattle at the following rate: 15mL per 100 animals per day. Do not exceed dose rate of 2mL per animal per day.

### Foliar application

Apply in a minimum of 200 litres water per hectare. Pasture Spraying: 1.5 to 2.5 litres per hectare ideally 4 days to 2 weeks prior to grazing. Repeat after 6 weeks as required. Treat approximately 1 hectare per 50 animals.

### Supplementary feed

Apply to cattle supplementary feed daily at the following rate: 15mL per 100 animals per day. Do not exceed 2mL per animal.

## STORAGE

Keep container closed. Store locked up in a cool place out of direct sunlight.

## DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product, packaging and waste at an approved landfill or equivalent facility in accordance with local, regional and national environmental regulations.

## FIRST AID

**IF SWALLOWED:** Rinse mouth. Give water to drink. If vomiting occurs give more water. Get medical advice/attention if you feel unwell.  
**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do, continue rinsing. If irritation persists, get medical advice/attention.  
**IF ON SKIN:** Take off contaminated clothing and wash skin with soap and water. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse.  
**IF INHALED:** If breathing is difficult, remove to fresh air and keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms, call a POISON CENTRE or doctor/physician.

If exposed or concerned, get medical advice.

**National Poisons Centre 24 Hour Phone:  
0800 POISON (0800 764 766)**

**DONAGHYS**  
PERFORMANCE FIRST

MANUFACTURED AND SOLD BY:  
DONAGHYS LTD  
16 SHEFFIELD CRESCENT  
PO BOX 20 449, HAREWOOD, CHRISTCHURCH  
FREEPHONE: 0800 942 006 WWW.DONAGHYS.COM



ZPLAB795 FEB25 PN-038860-1 VER2