

WARNING. FOR ANIMAL TREATMENT ONLY. KEEP OUT OF REACH OF CHILDREN.

DONAGHYS

Selenium 5

Dairy cattle and sheep mineral supplement



Selenium is a vital component of antioxidants that assist in livestock performance. Overcoming low Selenium levels can assist with:

- General good thrift
- Improved milk production
- Improved disease resistance
- Improved fertility

5mg/ml

DONAGHYS
PERFORMANCE FIRST



Donaghys Selenium 5

Help your cows and sheep achieve a higher level of performance with Selenium 5, the dairy cow and sheep liquid mineral supplement!

READ LABEL BEFORE USE

Keep out of reach of children. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid breathing mist/vapors/spray. In case of inadequate ventilation wear respiratory protection. Wear protective gloves/protective clothing/eye protection/face protection. Avoid release to the environment. Use only outdoors or in a well-ventilated area. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required.

DIRECTIONS

Only for use when there is a confirmed Selenium deficiency. Ready to use liquid no dilution required.

DONAGHYS SELENIUM 5 DOSE RATE

DO NOT EXCEED THE STATED DOSE RATE AND FREQUENCY

1ml will provide 5mg of Selenium.

Through water supply

Can be added to drinking water via dispenser. Administer daily to sheep and cattle at rates not exceeding the following:
0.02ml per 10kg of liveweight per day.
0.1ml per 50kg of liveweight per day.
0.2ml per 100kg liveweight per day.
1ml per 500kg of liveweight per day.

Oral drenching

Administer orally to sheep and cattle at rates not exceeding the following:
0.2ml per 10kg of liveweight once every 3 weeks.
1ml per 50kg of liveweight once every 3 weeks.
2ml per 100kg of liveweight once every 3 weeks.
10ml per 500kg of liveweight once every 3 weeks.

Do not administer more than once every 3 week period. Do not repeat drench more than 7 times a year.

Supplementary feed

Apply to supplementary feed daily at rates not exceeding the following:
0.02ml per 10kg of liveweight per day.
0.1ml per 50kg of liveweight per day.
0.2ml per 100kg of liveweight per day.
1ml per 500kg of liveweight per day.

CAUTION

Do not use at the same time as any other selenised fertiliser, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian. Do not repeat 3 weekly drenching more than 7 times per year. Cows pass on Selenium efficiently to the foetus. Pre-calving and post-calving are key times to supplement to ensure good level in both mother and calf but Selenium is beneficial year round.

STORAGE

Store locked up, in a cool place out of direct sunlight.

FIRST AID

IF SWALLOWED: Rinse mouth. Call a POISON CENTER or doctor/physician if you feel unwell. If medical advice is needed, have product container or label at hand.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

IF ON SKIN: Wash with plenty of soap and water. Wash contaminated clothing before reuse. Call a POISON CENTER or doctor/physician if you feel unwell. If medical advice is needed, have product container or label at hand.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing. Call a POISON CENTER or doctor/physician if you feel unwell. If medical advice is needed, have product container or label at hand.

IF exposed or concerned:
Get medical advice/attention.

**National Poisons Centre 24 Hour Phone:
0800 POISON (0800 764 766)**

DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product, packaging and waste at an approved landfill or equivalent facility in accordance with local, regional and national environmental regulations

HAZARDOUS SUBSTANCES CLASSIFICATION GHS7

Acute oral toxicity Category 4
Acute dermal toxicity Category 4
Acute inhalation toxicity Category 4
Germ cell mutagenicity Category 2
Specific target organ toxicity – single exposure Category 2
Specific target organ toxicity – repeated exposure Category 2
Hazardous to the aquatic environment chronic Category 3
Hazardous to soil organisms
Hazardous to terrestrial vertebrates

ZPLAB793 MAY24 PN-033832-1 VER2

DONAGHYS
PERFORMANCE FIRST

MANUFACTURED AND SOLD BY:
DONAGHYS LIMITED
16 SHEFFIELD CRESCENT
PO BOX 20 449, HAREWOOD, CHRISTCHURCH
FREEPHONE: 0800 942 006 WWW.DONAGHYS.COM

