

DONAGHYS

TRON

A SELECTIVE POST EMERGENT HERBICIDE FOR THE CONTROL OF CERTAIN WEEDS IN FODDER BEET, RED BEET, SUGAR BEET AND MANGOLDS.

10
Litre

Contains:

700 g/L Metamitron in the form of a suspension concentrate

TOXIC/ECOTOXIC
KEEP OUT OF REACH
OF CHILDREN

WARNING: TOXIC/ ECOTOXIC

Harmful if swallowed. Very toxic to aquatic life with long lasting effects. Very toxic to many plant species. Harmful to terrestrial vertebrates.

PRECAUTIONS:

APPROVED HANDLER: This product must be under control of an approved handler when used.

STORAGE: Keep out of reach of children. Read label before use. Store in original container away from feed or foodstuffs. Specific storage requirements apply based on quantities of product stored (Refer to Safety Note). When stored appropriately this product should not show any significant degradation for two years from the date of manufacture. Contact supplier for further instructions if product intended to be used is older than this.

PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. Wear protective overalls, boots, gloves eye/ face protection when mixing, loading or applying this product. An organic and particular matter respirator is required when using this product. Wash hands thoroughly after handling. Protective clothing has to be cleaned daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

USE: Apply with well-maintained and calibrated equipment. Apply at no more than 6L of product/ha and no more than three times per year.

DISPOSAL: Ensure unwanted product used by another operator is strictly in accordance with label directions. Triple rinse container and add residue to spray tank.

Donaghys is a member of Agrecovery Container Recycling programme. To recycle this container, please triple rinse and drop off to your local Agrecovery collection site. For details on how to correctly prepare your container and collection sites please visit the Agrecovery website www.agrecovery.co.nz

ENVIRONMENTAL: Use this product with care. Do not contaminate aquatic environments with product, spray or empty container. Do not apply directly into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Do not allow spray drift outside the target area to occur.

FIRST AID:

If medical advice is needed, have product container, label at hand. If swallowed do not induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. Rinse mouth.

If in eyes, wash with plenty of water for several minutes.

If on skin or hair, remove contaminated clothing and wash with plenty of soap and water. Call a POISON CENTER or doctor/ physician if you feel unwell. Wash contaminated clothing before reuse.

If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

SPILLS: Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways. Wear appropriate protective clothing (see Personal Protection section) to clean up spills. Absorb spills with an inert material (such as earth, sand or synthetic absorbent substance) and place in waste containers. Wash area with water and alkaline detergent then absorb any remaining liquid with further inert material. Dispose of the waste safely in an approved landfill.

SAFETY DATA SHEET: A Safety Data Sheet for this product is available from Donaghys Limited on request.

GENERAL INFORMATION:

Donaghys TRON is a selective post-emergence herbicide for the control of various weeds in Fodder beet, Red beet, Sugar beet and Mangolds. Uptake is predominantly via the roots, with some uptake via the leaves. Some residual activity may occur. For best results apply to small weeds.

Weeds controlled:

Susceptible: Amaranthus spp., annual poa, buttercups (seedling), chickweed, docks (seedling), fathen, galinsoga, groundsel, hawksbeard, henbit, matricaria spp. (stinking mayweed and chamomiles), nettles, pennycress, scotch thistle (seedling), shepherds purse, sow thistle, speedwells, spurrey, twin cress, white clover, wild portulaca, yarrow (seedling).

Moderately susceptible: Black nightshade, field pansy, fumitory, hedge mustard, scarlet pimpernel, thorn apple, wireweed.

Non-susceptible: Californian thistle, cleavers, cornbind, grasses (most), wild oats.

DIRECTIONS FOR USE:

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

CROP	RATE	CRITICAL COMMENTS
Fodder Beet, Red Beet, Sugar Beet, Mangolds	6L/ha in 200-300L/ha water	Apply early post emergence, when weeds are small, and the crop has fully expanded cotyledons but before the second true leaf stage.

APPLICATION METHOD:

Boom Spraying: Apply using conventional spray equipment with thorough agitation. Nozzles should produce a fine mist. Apply the required rate in 200-300L/ha water.

MIXING:

Shake well before use. Add the required quantity of Donaghys TRON to the partly filled with water spray tank, top up with water. Maintain agitation through filling and spraying.

RESISTANCE MANAGEMENT:

GROUP	5	HERBICIDE
-------	---	-----------

Donaghys TRON contains Group 5 herbicide belonging to the triazinone chemical family and continuous use of herbicides in this group may lead to a build-up of weeds resistant to it. To prevent or delay resistance, alternate with a herbicide from another group if possible and adhere to label instructions. Refer to <http://resistance.nzpps.org/index.php> for alternative modes of actions and details of resistance management strategies (if any) for the crops listed on the label.

IMPORTANT NOTICE - CONDITIONS OF SALE

ACCEPTANCE AND USE OF THIS PRODUCT WILL BE DEEMED TO BE ACCEPTANCE OF THESE CONDITIONS OF SALE.

Results from the use of this product may be affected by factors beyond Donaghys control. These factors include but are not limited to mixing, use, time of application, weather, crop, crop lifecycle and resistance to the active ingredient, even though the product is reasonably fit for its stated purpose and label directions have been followed. Donaghys will replace any defective product free of charge, but to the extent permitted by law all other warranties, representations, conditions or obligations whether imposed or implied by law or otherwise, and the manufacturer's and supplier's liability for any indirect special or consequential losses or damages of any kind arising out of the supply and/or use of this product, including where it arises out of negligence, are expressly excluded. **IN ANY EVENT, TO THE EXTENT PERMITTED BY LAW, THE MANUFACTURER'S AND SUPPLIER'S LIABILITY ARISING OUT OF THE SUPPLY AND OR USE OF THIS PRODUCT IS LIMITED TO THE AMOUNT PAID FOR THE SPECIFIC PRODUCT.**

In a transport emergency call 111; Police or Fire Brigade. For 24 hours specialist advice in an emergency, call 0800 CHEMCALL (0800 243 622). If additional safety information is required, refer to the Safety Data Sheet (SDS). For a copy of the SDS phone Donaghys Limited (0800 942 006).

HAZARD CLASSIFICATIONS: 6.1B, 6.1D, 9.1A, 9.2A, 9.3C

Acute oral toxicity Category 4

Hazardous to the aquatic environment acute Category 1

Hazardous to the aquatic environment chronic Category 1

Hazardous to soil organisms

Hazardous to terrestrial organisms

Registered pursuant to the ACVM Act 1997, No.P009587

See www.foodsafety.govt.nz for registration conditions.

Approved Pursuant to the HSNO Act 1996, No. HSR100753

See www.epa.govt.nz for approval conditions



UN number 3082
Packaging Group III
Hazchem 3Z
**Environmentally Hazardous Substance,
Liquid, N.O.S. (Contains Metamitron 70%)**



DONAGHY'S
PERFORMANCE FIRST

REGISTERED AND MANUFACTURED BY DONAGHY'S LTD
16 SHEFFIELD CRESCENT, PO BOX 20 449, HAREWOOD, CHRISTCHURCH
PHONE 0800 942 006 www.donaghys.com

ZPLAB761 MAY22 PN-030236-1 VER3