

**WARNING**

KEEP OUT OF REACH OF CHILDREN  
READ ENTIRE LABEL BEFORE USE  
FOR ANIMAL TREATMENT ONLY

**DONAGHYS**

# Epripor X



**BROAD SPECTRUM  
POUR-ON ENDECTOCIDE  
FOR CATTLE**

For the Treatment and Control of all  
Gastrointestinal Roundworms,  
Lungworms and Sucking Lice  
in Cattle

ACTIVE CONSTITUENT  
Each ml contains 5 mg eprinomectin

NET CONTENTS  
**20 Litres**

**DONAGHYS**  
PERFORMANCE FIRST

DONAGHYS

# Epripor X BROAD SPECTRUM POUR ON ENDECTOCIDE FOR CATTLE

**READ THE ENTIRE LABEL BEFORE USE**

May cause skin and eye irritation. May damage fertility or the unborn child from repeated oral exposure. Danger - May cause harm to breast fed children from repeated oral exposure.

**Handling Precautions:** Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. Avoid contact with skin and eyes. When handling wear waterproof gloves and eye protection. Do not breathe mist/vapours/spray. Wash hands and face thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid contact during pregnancy/while nursing.

**First Aid: IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

**IF ON SKIN:** Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before re-use. IF exposed or concerned: Get medical advice/attention. For advice, contact the National Poisons Centre - 0800 POISON (0800 764 766) or a doctor immediately.

**Environmental Protection:** Very toxic to aquatic organisms with long lasting effects. Harmful to the soil environment. Harmful to terrestrial vertebrates. Very toxic to terrestrial invertebrates. Avoid contamination of any water supply with product or empty container. Avoid release to the environment. Collect spillage.

**Emergency Response:** In case of spillage wear appropriate protective clothing. Prevent material from entering waterways. Absorb spilled products with inert material and place in waste containers. Sweep up solids without creating a dust. Wash area with further inert material. Dispose of waste safely.

Also contains NMP: 10-20% w/v

Epripor X is a broad spectrum endectocide. It contains eprinomectin, a member of the macrocyclic lactone (avermectin/milbemycin) family. It delivers internal parasite control in one low volume application. It's convenience, broad-spectrum efficacy, zero milk withholding period, dose, and wide margin of safety make it a suitable product for parasite control in beef and dairy cattle.

**INDICATIONS:** For the treatment and control of internal parasites, lungworm and sucking lice in all types of cattle.

**Gastrointestinal worms:** *Ostertagia ostertagi* (including inhibited immature), *Ostertagia lyrata*, *Haemonchus* spp., *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Oesophagostomum radiatum*, *Nematodirus helvetianus*, *Bunostomum phlebotomum*, *Trichuris* spp. and *Cooperia* spp.

**Lungworm:** Adult and immature *Dictyocaulus viviparus*.

**External Parasites:**

Sucking Lice: *Linognathus vituli*, *Haematopinus eurysternus* and *Solenopotes capillatus*.

Resistance can develop to any anthelmintic. Macrocyclic lactone resistant *Cooperia* are becoming widespread in New Zealand, therefore efficacy against *Cooperia* may be reduced.

Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

**DIRECTIONS FOR USE:**


By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

For external use only. Shake well before use. Correct drenching technique should be used. Draft animals into lines of similar bodyweight. Choose dose according to the weight of the heaviest animal in each line. Check accuracy of drench guns regularly. Do NOT underdose.

Administer topically along the midline of the back, from the withers to the base of the tail, using standard pour-on equipment. Do not use if rainfall is pending.

**DOSE RATE:** 1ml per 10kg bodyweight, equivalent to 500µg eprinomectin per kg bodyweight. Epripor X is not recommended for use on calves under 100kg bodyweight.

**DRUM TAP ENCLOSED.**  
Fit to internal thread after removing small drum cap.



Cut along dotted line

Bodyweight (kg)	Dose Volume (ml)	Doses per 20L Pack
101-120	12	1666
121-140	14	1428
141-160	16	1250
161-180	18	1111
181-200	20	1000
201-250	25	800
251-300	30	666
301-350	35	571
351-400	40	500
401-450	45	444
451-500	50	400
501-550	55	363
551-600	60	333
601-650	65	307
Over 650kg add 5ml for every 50kg increase in bodyweight.		

**WARNINGS AND CONTRAINDICATIONS:**

Do not use on animals which are severely stressed or in ill health. Care should be taken in handling pregnant animals. Studies have demonstrated a wide safety margin. Three times the recommended dose had no adverse effect on breeding performance of cows or bulls. Five times the recommended dose had no other adverse effects on cattle.

**WITHHOLDING PERIODS:**

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**MEAT** - There is no meat withholding period for cattle

**Milk** - There is no milk withholding period for cattle

**Bobby Calves** - There is no withholding period for bobby calves born to cows treated with Epripor X.

**STORAGE:** Store locked up below 25°C (Room Temperature). Protect from direct sunlight. Keep container tightly closed when not in use.

**DISPOSAL:** Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill/facility. Triple rinse container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/facility. Do not use container for any other purpose.

Registered pursuant to the ACVM Act 1997, No A010357  
See www.foodsafety.govt.nz for registration conditions.

**DONAGHYS**

Registered to and distributed by Donaghys Ltd, 16 Sheffield Crescent, Harewood, Christchurch, New Zealand  
Freephone: 0800 942 006  
www.donaghys.com

DONAGHYS

# Epripor X

BROAD SPECTRUM POUR ON ENDECTOCIDE FOR CATTLE



UN3082 ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (Eprinomectin)



Batch:  
Expiry: