

**WARNING**

KEEP OUT OF REACH  
OF CHILDREN  
READ LABEL  
BEFORE USE  
FOR ANIMAL  
TREATMENT ONLY

**DONAGHYS**

# MoxiPor

## POUR ON



### LONG LASTING ENDECTOCIDE FOR CATTLE AND DEER

For the treatment and control of internal and external parasites of cattle (including lactating dairy cattle) and for the treatment and control of lungworm and roundworms of deer.

Improves milk production in lactating dairy cows and increases weight gains in growing beef and dairy cattle exposed to parasite challenge.

NET CONTENTS  
**20 Litres**

ACTIVE CONSTITUENT  
Moxidectin 5g/L  
Contains Aromatic Hydrocarbons 150g/L

**DONAGHYS**  
PERFORMANCE FIRST

DONAGHYS

# MoxiPor POUR ON LONG LASTING ENDECTOCIDE FOR CATTLE AND DEER

**READ LABEL CAREFULLY BEFORE USING** Read label carefully and obtain special instructions before use. Do not handle it until all safety precautions have been read and understood.

**HANDLING PRECAUTIONS** May cause mild skin irritation. Moxidectin may possibly affect development and/or reproduction, and may have effects on or via lactation. Do not breathe mist, vapours or spray. Avoid skin contact. Avoid contact during pregnancy/while nursing. Handle with care. The use of chemically resistant gloves is recommended during use. Wear protective gloves and use personal protective equipment, as required. Do not eat, drink or smoke when using this product and wash hands thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace and should be washed before reuse. Store locked up. Harmful if swallowed.

**FIRST AID** For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water.

**ENVIRONMENTAL PROTECTION** Very toxic to the aquatic environment. Avoid contamination of any water supply with product or empty container. Harmful to the soil environment. Toxic to terrestrial invertebrates. Avoid release to the environment. Harmful to terrestrial vertebrates.

**EMERGENCY RESPONSE** In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely.

**DISPOSAL** Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Triple rinse the empty container, crush or puncture and bury in a suitable landfill, or if appropriate, recycle. Do not use container for any other purpose

DONAGHYS

# MoxiPor POUR ON LONG LASTING ENDECTOCIDE FOR CATTLE AND DEER



**MOXIPOR POUR ON** contains moxidectin, a member of the milbemycin/ avermectin family. It is effective against sensitive strains of the following internal parasites:

**CATTLE Gastrointestinal Roundworms:** Adult and immature forms unless otherwise stated; *Haemonchus contortus* (barber's pole worm), *Ostertagia ostertagi* (*O. lyrata*) (small brown stomach worm including inhibited larvae), *Trichostrongylus axei* (stomach hair worm), *Trichostrongylus colubriformis* (black scour worm), *Trichostrongylus longispicularis*, *Cooperia oncophora* (small intestinal worm), *Cooperia punctata*, *Oesophagostomum radiatum* (nodule worm), *Bunostomum phlebotomum* (hookworm).

**Lungworm:** Adult and immature *Dictyocaulus viviparus*. **Lice:** *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*, *Bovicola bovis*.

**Mange mites:** *Chorioptes bovis*.

**DEER Lungworm:** *Dictyocaulus ekerti* (viviparus).

**Gastrointestinal roundworms:** *Haemonchus contortus*, *Ostertagia*-like species\*, *Trichostrongylus* spp, *Oesophagostomum venulosum*, *Chabertia ovina*

\**Ostertagia leptospicularis* and *Ostertagia*-like species including *Spiculoptera assymetrica*, *Spiculoptera spiculoptera* and *Skrjabinagia kolchida*.

**PERSISTENT ACTIVITY** MOXIPOR POUR ON, following a single application at the recommended dose rate, prevents reinfection of:

**CATTLE** with *Dictyocaulus viviparus*, *Oesophagostomum radiatum* and *Bunostomum phlebotomum* for 42 days, *Ostertagia ostertagi* (*O. lyrata*) for 35 days and *Haemonchus* spp and *Trichostrongylus axei* for 28 days, and

**DEER** with *Dictyocaulus ekerti* (viviparus), *Ostertagia*-like spp\*, *Oesophagostomum venulosum* and *Chabertia ovina* for 42 days.

\**Ostertagia leptospicularis* and *Ostertagia*-like species including *Spiculoptera assymetrica*, *Spiculoptera spiculoptera* and *Skrjabinagia kolchida*.

Resistance may develop to any anthelmintic. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance.



**DOSAGE AND ADMINISTRATION:**

Dose Rate: 1 mL/10 kg to give 0.5 mg moxidectin per kg liveweight.

Liveweight (kg)	Dose Volume (mL)	Doses per 20L Pack	Liveweight (kg)	Dose Volume (mL)	Doses per 20L Pack
50	5	4000	250	25	800
60	6	3333	300	30	666
70	7	2857	350	35	571
80	8	2500	400	40	500
90	9	2222	450	45	444
100	10	2000	500	50	400
110	11	1818	550	55	363
120	12	1666	600	60	333
130	13	1538	650	65	307
140	14	1428	700	70	285
150	15	1333	750	75	266
200	20	1000			

Registered pursuant to the ACVM Act 1997, No. A11390 See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions. Approved pursuant to the HSNO Act, No HSR001850 See <http://www.epa.govt.nz> for controls.

**DIRECTIONS FOR USE:**

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Ready for use with standard pour-on equipment. MOXIPOR POUR ON should be applied along the top of the back from the shoulders to the base of the tail. Check accuracy of the pour-on applicator regularly. A representative sample of animals should be weighed before treatment either with scales or a weighband. Choose dose according to the weight of the heaviest animals. Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight to avoid excessive overdosing.

**Rainfast:** The efficacy of MOXIPOR POUR ON is not adversely affected if applied when the MoxiPor hide is wet, or if rain occurs shortly after the application. However treatment of animals under these circumstances is not recommended practice.

**Non-irritant:** MOXIPOR POUR ON does not cause irritation to the skin or damage to the hide when used as directed.

**Ease of Use:** MOXIPOR POUR ON comes ready to use in applicator bottles, back packs or drums.

**Concurrent Use:** MOXIPOR POUR ON is safe to use concurrently with other treatments in cattle including mineral supplements, fluke treatments and vaccines. It should not be applied at the same time as any other pour-on product.

**Safety to Target Animals:** MOXIPOR POUR ON has a wide margin of safety when used at the recommended dose rate.

**WITHHOLDING PERIODS:**

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

- MEAT
- Cattle - Nil
- Deer - Nil
- Bobby Calves - Nil
- Milk - Nil

**STORAGE:** Store below 25°C and protect from light.

## DONAGHYS

Registered to:  
Horizon Agresources (NZ) Limited

Marketed and Distributed by:  
Donaghys Ltd  
16 Sheffield Crescent,  
Harewood, Christchurch, New Zealand  
Freephone: 0800 942 006  
[www.donaghys.co.nz](http://www.donaghys.co.nz)

UN3082  
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.  
(Contains Moxidectin 0.5%)  
Packaging Group III



Batch:  
Expiry: