

DONAGHYS

# ProAbamec

## POUR-ON FOR CATTLE

For the treatment and control of internal and external parasites of cattle.

**WARNING**  
**FLAMMABLE LIQUID**  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

ACTIVE INGREDIENTS  
**Abamectin** 10mg/ml  
Contains >941g/L  
Diethylene glycol monobutyl ether

NET CONTENTS  
**5 Litre**

**DONAGHYS**  
PERFORMANCE FIRST

**DONAGHYS**  
**ProAbamec**

READ ENTIRE LABEL BEFORE USE

**WARNING: Dangerous to the environment.**

**HANDLING PRECAUTIONS** Keep away from heat/sparks/open flames/hot surfaces. Wear protective gloves/protective clothing/eye protection. Keep closed when not in use. May be harmful if swallowed, inhaled or absorbed through the skin.

May cause mild skin and eye irritation. Repeated exposure may cause skin allergy. Abamectin may possibly affect development and/or reproduction and may have effects on/ or via lactation. Diethylene glycol monobutyl ether may possibly affect kidney if swallowed and liver if inhaled. Wash hands and exposed skin before meals and after use. Do not eat, drink or smoke when using this product. Do not inhale vapour. Avoid skin and eye contact. Very toxic to aquatic organisms and terrestrial invertebrates. Toxic to the soil environment. Harmful to terrestrial vertebrates. Avoid contamination of any water supply with product or empty container. Avoid release to the environment.

**FIRST AID** If swallowed do NOT induce vomiting. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If splashed in eyes wash out immediately with water. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

**EMERGENCY RESPONSE** In case of spillage wear appropriate protective clothing. Prevent material from entering waterways and absorb spills with inert material and place in waste container. Wash area and absorb with further inert material.

**INDICATIONS** Donaghys ProAbamec Pour-On for Cattle, when used at the recommended dose level of 500 micrograms Abamectin per kg liveweight, effectively controls the following parasites of cattle:

**GASTRO-INTestinal ROUNDWORMS** Adult and immature *Ostertagia* spp (including inhibited immatures), *Haemonchus* spp, *Trichostrongylus* spp, *Cooperia* spp, *Bunostomum* spp, *Oesophagostomum* spp, and adults of *Chabertia* spp, *Nematodirus* spp, *Trichuris* spp, and *Strongyloides* spp.

**LUNGWORMS** Adult and immature *Dictyocaulus viviparus*.

**BITING AND SUCKING LICE** *Bovicola bovis*, *Linognathus vituli*.

**DIRECTIONS FOR USE** By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Donaghys ProAbamec Pour-On for Cattle contains 10mg/ml Abamectin, a member of the macrocyclic lactone group of compounds. It is effective against internal and external parasites sensitive to this family. Resistance can develop to any anthelmintic. Check the effectiveness of anthelmintics at least annually. Follow recommended parasite management practices to reduce the potential for resistance to develop. Check accuracy of pour-on applicator regularly. A representative sample of animal should be weighed before treatment. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc). Do not under-dose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Ready to use

with standard pour-on equipment. Apply along the backline from shoulders to tail.

Donaghys ProAbamec Pour-On is rainfast, even if rain falls 2 hours or more after treatment.

**DOSAGE** It is strongly recommended not to use this product in cattle under four months of age and/or 100kg live weight. The dose rate for cattle is 500µg Abamectin per kg bodyweight, which is provided by 1ml Donaghys ProAbamec Pour-On for Cattle per 20kg bodyweight, or according to the table:

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

### WITHHOLDING PERIODS

**MILK** Milk intended for sale for human consumption must be discarded during treatment and for **35 days** following the last treatment.

**MEAT** Cattle producing meat and offal for human consumption must not be sold for slaughter during or within **35 days** of the last treatment.

**STORAGE** Store below 25°C. Protect from sunlight and keep it well closed.

**DISPOSAL** Preferably dispose of product by use. Otherwise dispose of product, packaging and waste at an approved landfill or equivalent facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate recycle.

Registered pursuant to the ACVM Act 1997 No.A010875.  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.  
Approved pursuant to the HSNO Act 1996, approval code HSR100682.  
See [www.epa.govt.nz](http://www.epa.govt.nz) for controls.

### REGISTERED TO AND DISTRIBUTED BY

Donaghys Ltd, 16 Sheffield Crescent, Harewood, Christchurch  
Freephone: 0800 942 006  
[www.donaghys.com](http://www.donaghys.com)

UN No.3082, Packaging Group III, HazChem 3Z  
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (Contains Abamectin)

DOSE RATE		
Live-weight (kg)	Dose (mL)	5l Pack Treats
101-150kg	7.5mL	666
151-200kg	10mL	500
201-250kg	12.5mL	400
251-300kg	15mL	333
301-350kg	17.5mL	285
351-400kg	20mL	250
401-450kg	22.5mL	222
451-500kg	25mL	200
501-550kg	27.5mL	181
551-600kg	30mL	166
601-650kg	32.5mL	153

Heavier animals (e.g. mature bulls) should receive an additional 1ml for each 20kg over 650kg.



Batch:  
Expiry:

2PLAB2 SEP24 PN 03M0751 V055

