

WARNING

KEEP OUT OF REACH
OF CHILDREN
READ LABEL BEFORE USE
FOR ANIMAL
TREATMENT ONLY

DONAGHYS

Saturn[®]

 **POUR-ON FOR CATTLE**

For the treatment and control
of internal parasites, including
endectocide resistant strains,
lungworm and sucking
lice in cattle.

NET CONTENTS
2.5 Litres

DONAGHYS
PERFORMANCE FIRST

**ACTIVE
CONSTITUENTS**

Each mL contains 10mg abamectin
and 200mg levamisole base. Also contains
35-45%w/v glycerol formal stabilized
(as the vehicle).

WARNING

Dangerous to the Environment
READ ENTIRE LABEL BEFORE USE

May be harmful if swallowed, inhaled or absorbed through the skin. May cause an allergic skin reaction. Suspected of causing heritable mutations/genetic defects. Suspected of damaging fertility or the unborn child from repeated exposure. May cause harm to breast-fed children from repeated exposure.
Danger - Presumed to/may cause organ damage from repeated exposure at high doses (levamisole may affect the blood and haematopoietic system).

HANDLING PRECAUTIONS Avoid skin contact. When handling wear chemically-resistant gloves. Remove protective clothing and wash hands and face thoroughly before meals and after work. Do not eat, drink or smoke when using this product.

FIRST AID If swallowed, do NOT induce vomiting. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

ENVIRONMENTAL PROTECTION Very toxic to aquatic organisms with long lasting effects. Harmful to the soil environment. Toxic to terrestrial vertebrates. Very toxic to terrestrial invertebrates. Avoid contamination of any water supply with product or empty container. Avoid release to the environment. Collect spillage.

EMERGENCY RESPONSE For specialist advice in an emergency call 0800 734 607 (24 hours).

SATURN POUR-ON is a broad spectrum endectocide. It contains abamectin, a member of the macrocyclic lactone (avermectin/milbemycin) family. It also contains levamisole base, a member of the imidazothiazole class of anthelmintics, with efficacy against internal parasites sensitive to this family, especially lungworm and Cooperia.

INDICATIONS For the treatment and control of mature and immature stages of internal parasites, including endectocide-resistant strains, and lungworm in all types of cattle.

Gastrointestinal worms: *Ostertagia ostertagi**, *Ostertagia lyrata**, *Cooperia oncophora**, *Cooperia sumabada**, *Cooperia punctata**, *Trichostrongylus longispicularis*, *Trichostrongylus axei*, and mature *Trichuris* (*includes inhibited L4 stage larvae).

SATURN POUR-ON is also highly effective in the treatment and control of mature and immature strains of *Cooperia* spp. resistant to the endectocides (including eprinomectin and doramectin). Levamisole is also very active against susceptible benzimidazole-resistant strains.

Lungworm: Adult and immature *Dictyoacaulus viviparus*.

External Parasites: For the treatment and control of sucking lice in cattle, including: *Linognathus vituli*, *Haematopinus eurystermus*, *Solenopotes capillatus*.

SATURN POUR-ON is effective against all internal parasites sensitive to either or both active families. Resistance can develop to any anthelmintic. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

DIRECTIONS FOR USE By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

For external use only. Shake well before use. Correct drenching technique should be used. A representative sample of animals should be weighed before treatment. Draft animals into lines of similar bodyweight. Choose dose according to weight of the heaviest animal in each line. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight to avoid excessive overdosing. Check accuracy of drench guns regularly. Do NOT underdose.

Administer topically onto the skin along the midline of the back, from the withers to the base of the tail, using standard pour-on equipment.

DOSE 1mL per 20kg bodyweight, equivalent to 500µg abamectin and 10mg levamisole per kg bodyweight. Do not use on calves under 60kg bodyweight.

WARNINGS AND CONTRAINDICATIONS

Do not use on animals which are severely stressed or in ill health. Care should be taken in handling pregnant animals. Do not exceed recommended dose rates. Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity, so estimate liveweights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between levamisole and organophosphate dips.

WITHHOLDING PERIODS It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MEAT: Cattle producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 42 days of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 42 days following the last treatment.

Bodyweight	Dose	Bodyweight	Dose
61 – 80kg	4.0mL	361 – 380kg	19.0mL
81 – 100kg	5.0mL	381 – 400kg	20.0mL
101 – 120kg	6.0mL	401 – 420kg	21.0mL
121 – 140kg	7.0mL	421 – 440kg	22.0mL
141 – 160kg	8.0mL	441 – 460kg	23.0mL
161 – 180kg	9.0mL	461 – 480kg	24.0mL
181 – 200kg	10.0mL	481 – 500kg	25.0mL
201 – 220kg	11.0mL	501 – 520kg	26.0mL
221 – 240kg	12.0mL	521 – 540kg	27.0mL
241 – 260kg	13.0mL	541 – 560kg	28.0mL
261 – 280kg	14.0mL	561 – 580kg	29.0mL
281 – 300kg	15.0mL	581 – 600kg	30.0mL
301 – 320kg	16.0mL	601 – 620kg	31.0mL
321 – 340kg	17.0mL	621 – 640kg	32.0mL
341 – 360kg	18.0mL	641 – 660kg	33.0mL

Over 660kg add 1mL for every 20kg increase in bodyweight.

STORAGE Store locked up below 25°C (room temperature). Protect from light. Keep container tightly closed when not in use.

DISPOSAL Treat the product so that it is no longer toxic. Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Triple rinse empty container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/facility. Do not use container for any other purpose.

Registered pursuant to the ACVM Act 1997, No A010206
See www.foodsaftey.govt.nz for registration conditions.

Donaghys Ltd, 16 Sheffield Crescent,
Harewood, Christchurch, New Zealand
Freephone: 0800 942 006
www.donaghys.com

UN 3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE,
LIQUID, N.O.S., (1% abamectin)

