

DONAGHYS

StrikeOut



SPRAY-ON

For the preventative control of blowfly strike on sheep.

NET CONTENTS

10 Litres

ACTIVE INGREDIENT
Cyromazine 60g/L

WARNING

FOR ANIMAL
TREATMENT ONLY
READ THE ENTIRE
LABEL BEFORE USE

DONAGHYS
PERFORMANCE FIRST

READ THE ENTIRE LABEL BEFORE USE

INDICATIONS StrikeOut Spray-On is an easy to use spray which will give up to 6 weeks residual protection against flystrike caused by common strike flies including the Australian blowfly (*Lucilia cuprina*).

StrikeOut Spray-On may be applied to all breeds of sheep, and is suitable for use between 4 and 12 weeks after shearing.

LIMITATIONS The period of protection may be reduced if StrikeOut Spray-On is applied incorrectly, if there is excessive fleece soiling or if wool infection such as mycotic dermatitis (fleece rot) is present. The protection period may also be reduced by prolonged heavy rain following treatment.

DIRECTIONS FOR USE *By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.*

DO NOT DILUTE. DO NOT mix with any other product. Apply directly from the container using a suitable clean applicator gun. The product should be sprayed on in bands, with the gun held 20cm above the wool during application. StrikeOut Spray-On should be applied onto those areas where maximum protection against flystrike is required.



StrikeOut Spray-On applied between the rump and shoulders should be sprayed on in even bands as shown (*see diagram*).


The product sprayed onto the rump should be applied in an arc over the area shown (*see diagram*).

The dose rate and number of bands to be applied are shown in the table. Note: The dose volume applied should be based on the weight of the heaviest sheep in the mob.

APPLICATION AND DOSE RATES OF STRIKEOUT SPRAY-ON

For the protection of all the commonly struck areas of the sheep:

	2 bands sheep between 11-20kg		3 bands sheep between 21-40kg		4 bands sheep between 41-60kg		5 bands
Liveweight (kg)	11-15	16-20	21-30	31-40	41-50	51-60	61-70
No. of bands	2	2	3	3	4	4	5
Volume per band (mL)	7.5	10	10	13.3	12.5	15	14
Total vol applied (mL)	15	20	30	40	50	60	70
Sheep treated per 10L	666	500	333	250	200	166	142



For prevention of flystrike at docking: Apply 5 – 10mL over the crutch, ensuring that sufficient StrikeOut Spray-On is applied to cover the area to be protected (*see diagram*).

For the prevention of poll strike: Rams and sheep susceptible to poll strike should be treated. Apply at least 5mL StrikeOut Spray-On to the areas requiring protection, around the poll and horns (*see diagram*).

WITHHOLDING PERIODS *It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.*

MILK: Milk intended for sale for human consumption must be discarded during treatment and for **35 days** following the last treatment.

MEAT: Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within **7 days** of the last treatment.

WOOL RESIDUES: Ensure the time between dipping and subsequent shearing is at least **2 months**, except in the case of spot treatment.

STORAGE Store below 30°C (room temperature), in original containers, tightly closed in a dry, cool place. DO NOT store for prolonged periods in sunlight. DO NOT reuse the container.

HANDLING PRECAUTIONS:

May cause skin and eye irritation. May possibly affect the blood and the hematopoietic system. Avoid skin and eye contact. When using the product wear rubber gloves. Wash hands after use.

KEEP OUT OF REACH OF CHILDREN

FIRST AID If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If splashed in eyes wash out immediately with water. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

ENVIRONMENTAL PROTECTION Harmful to aquatic organisms. Avoid contamination of any water supply with product or empty container.

EMERGENCY RESPONSE In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material.

DISPOSAL Preferably dispose of product by use. Otherwise dispose of product, packaging and waste at an approved landfill or equivalent facility.

Registered to: Abbey Animal Health NZ Ltd
68 Russell Street, Westport 7825, New Zealand

Marketed and Distributed by: Donaghys Ltd
16 Sheffield Cres, Harewood, PO Box 20 449, Christchurch 8543, New Zealand
Freephone: 0800 942 006 www.donaghys.com

Registered pursuant to the ACVM Act 1997 No. A9952.
See www.foodsafety.govt.nz for registration conditions.



UN 3082
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CYROMAZINE)
PG III