




Donaghys GibbSTART MAX

GibbSTART MAX stimulates production of winter dormant grass-dominant pastures for high intensity grazing such as dairy pasture or sheep lambing paddocks.

Active Ingredient: Contains 200g/L Gibberellic acid in the form of a soluble concentrate.

- Stimulates extra pasture production when extra feed is needed in late Autumn and early Spring when pasture growth is limited by cooler soil temperatures.
- Liquid formulation eliminates dissolving and dust issues.
- Easy to mix and measure.
- Can be tank mixed with Donaghys selective herbicides 2,4-D Amine and 2,4-D Ester 680.
- Compatible with Donaghys N-Boost® and dissolved urea fertiliser.
- Available in 1L – enough to cover 25 hectares!



WARNING:

As with all gibberellin applications, the use of GibbSTART MAX can cause rapid pasture growth utilizing stored energy reserves of the plants. There may be a subsequent lag in growth. Such a lag becomes more likely if repeat applications are made without time for recovery of reserves. Quick repeat applications are likely to give diminishing returns. We recommend one-off strategic applications to address a feed deficit.

DO NOT apply to pasture more than four times in a calendar year. For best results, apply when there is sufficient soil moisture and soil nutrient availability to support rapid plant growth. Soil temperature should be less than 13°C for optimum response.

Some weeds may become more palatable to stock so avoid application to pastures containing high populations of toxic weeds.

Pastures including over sown grasses should be at least 1 year old prior to a GibbSTART MAX application.

Pasture Management: Growth stimulation is usually seen within 7 days of application. Do not graze within 2 weeks of an application as pasture quality is affected by gibberellin application. Grazing should recommence within 28 days of application to ensure the extra pasture growth is utilized and grass does not become rank.

DIRECTIONS FOR USE:

- Apply at 40ml/ha in total volume of water of 100 litres/ha
- Apply GibbSTART MAX 3 to 5 days after grazing when the pasture has started to grow again.
- Do not apply if rain is expected within 2 hours of application
- For best results GibbSTART MAX should be applied with a non-ionic surfactant such as Donaghys Wetter Sticker

Code	Product	Unit	Qty per Unit	Barcode
AGIBM001	DONAGHYS GIBBSTART MAX 1L	EACH	1	9400579552794

Registered pursuant to the ACVM Act 1997 No.P8774.

Refer to product label for complete usage, safety and product information before use.