

DONAGHYS

HiMin

# Venator<sup>®</sup>

LOW DOSE  
DOUBLE  
ACTIVE

## ORAL DRENCH FOR SHEEP AND CATTLE

For the treatment and control of all major gastrointestinal parasites in sheep and cattle, including strains with single resistant to benzimidazole (BZ), levamisole (LEV) and avermectin/milbemycin (ML) action families or to combinations of ML + BZ, or BZ + LEV action families.

Also provides supplementary source of cobalt and **selenium**.

NET CONTENTS  
**10 Litres**

**DANGER**  
READ LABEL BEFORE USE  
KEEP OUT OF THE  
REACH OF CHILDREN  
FOR ANIMAL  
TREATMENT ONLY

**ACTIVE  
INGREDIENTS**  
Abamectin 2g/L; Levamisole  
Hydrochloride 80g/L  
Also contains: **Selenium** 5mg  
per 5mL and Cobalt 25mg per  
5mL

**DONAGHYS**  
PERFORMANCE FIRST

DONAGHYS

HiMin

# Venator<sup>®</sup>

LOW DOSE  
DOUBLE  
ACTIVE

## ORAL DRENCH FOR SHEEP AND CATTLE

For the treatment and control of all major gastrointestinal parasites in sheep and cattle, including strains with single resistant to benzimidazole (BZ), levamisole (LEV) and avermectin/milbemycin (ML) action families or to combinations of ML + BZ, or BZ + LEV action families.

Also provides supplementary source of cobalt and **selenium**.

NET CONTENTS  
**10 Litres**

**DANGER**  
READ LABEL BEFORE USE  
KEEP OUT OF THE  
REACH OF CHILDREN  
FOR ANIMAL  
TREATMENT ONLY

**ACTIVE  
INGREDIENTS**  
Abamectin 2g/L; Levamisole  
Hydrochloride 80g/L  
Also contains: **Selenium** 5mg  
per 5mL and Cobalt 25mg per  
5mL

**DONAGHYS**  
PERFORMANCE FIRST

**READ ENTIRE LABEL BEFORE USE**

**INDICATIONS** Donaghys Venator HiMin contains levamisole, a member of the imidazothiazole family, abamectin, a member of the macrocyclic lactone family. It is effective against gastrointestinal worms sensitive to one or two or both drench families. For the effective treatment and control of mature and immature stages of susceptible strains of the following gastrointestinal parasites (\* includes inhibited fourth stage larvae) and lungworm:

Roundworms	Genus	Sheep	Cattle
<b>Abomasum</b> Barber's Pole Worm*	<i>Haemonchus*</i>	<i>contortus</i>	
Small Brown Stomach Worm*	<i>Teladorsagia*</i> <i>Ostertagia*</i>	<i>circumcincta</i>	<i>ostertagi</i>
Stomach Hair Worm	<i>Trichostrongylus</i>	<i>axeii</i>	<i>axeii</i>
<b>Small Intestine</b> Black Scour Worm	<i>Trichostrongylus</i>	<i>colubriformis</i> <i>vitrinus</i>	<i>longispicularis</i>
Small Intestinal Worm*	<i>Cooperia</i>	<i>curticei</i>	<i>oncophora</i> <i>punctata</i>
Thin-Necked Intestinal Worm	<i>Nematodirus</i>	<i>filicollis</i> <i>spathiger</i> <i>abnormalis</i>	<i>helvetianus</i>
Hookworm	<i>Bunostomum</i> <i>Capillaria</i> <i>Strongyloides</i>	<i>trigonocephalum</i> <i>papillosus</i>	<i>phlebotomum</i> <i>bovis</i>
<b>Large Intestine</b> Large Bowel Worm	<i>Oesophagostomum</i>	<i>venulosum</i>	<i>radiatum</i>
Large-Mouthed Bowel Worm	<i>Chabertia</i>	<i>ovina</i>	
Whipworm	<i>Trichuris</i>	<i>ovis</i>	
<b>Lungworm</b>			
Large Lungworm	<i>Dictyocaulus</i>	<i>filaria</i>	<i>viviparus</i>

**Anthelmintic resistant strains:** For the effective treatment and control of susceptible strains of gastrointestinal worms with single resistance to benzimidazole (BZ), levamisole (LEV) and avermectin/milbemycin (ML) action families or to combinations of ML + BZ, or BZ + LEV action families.

**DOSE RATE**

**SHEEP 1mL per 10kg Liveweight**

Liveweight	Dose	Doses per 10L pack
Less than 20 kg	DO NOT USE	
20-30kg	3mL	3333
31-40kg	4mL	2500
41-50kg	5mL	2000
51-60kg	6mL	1667
61-70kg	7mL	1429

Over 70kg add 1ml for every 10kg liveweight

**CATTLE 1mL per 10kg Liveweight**

Liveweight	Dose	Doses per 10L pack
Less than 100 kg	DO NOT USE	
101-150kg	15mL	666
151-200kg	20mL	500
201-250kg	25mL	400
251-300kg	30mL	333
301-350kg	35mL	285
351-400kg	40mL	250
401-450kg	45mL	222
451-500kg	50mL	200
501-550kg	55mL	181
551-600kg	60mL	166
601-650kg	65mL	153

Over 650kg add 1ml for every 10kg liveweight

1ml per 10kg bodyweight equivalent to levamisole hydrochloride 8mg, abamectin 0.2mg and selenium 0.1mg per kg bodyweight.

**DIRECTIONS FOR USE:** By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using this product other than as directed on the label.

**SHAKE WELL BEFORE USE.** For oral administration only. Ensure formulation remains evenly suspended during use. A representative sample of animals should be weighed before treatment. Draft animals into lines of similar bodyweight, then dose according to the weight of the heaviest animal in the line. When there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing. Do not under dose. Correct drenching technique should be used. Check accuracy of drench guns regularly. Rinse gun with water immediately after use.

Accuracy of drench guns should be checked before and during drenching. Ensure that the dosing technique is accurate and that each animal receives and swallows the full dose.

**WITHHOLDING PERIODS** It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: **Maximum Residue Levels for Agricultural Compounds**

**MEAT** **Sheep: 14 days** - Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 14 days of the last treatment.

**Cattle: 14 days** - Cattle producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 14 days of the last treatment.

**MILK** **35 days** - Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

**STORAGE** Store locked up below 25°C, out of direct sunlight.

**DISPOSAL** Preferably dispose of product by use. Otherwise dispose of product, packaging and waste at an approved landfill or equivalent facility in accordance with local, regional and national environmental regulations. Donaghys is a member of Agrecovery Container Recycling programme. For details on how to correctly prepare your container and collection sites please visit [www.agrecovery.co.nz](http://www.agrecovery.co.nz).



www.agrecovery.co.nz



Harmful if swallowed. Causes mild skin irritation. May cause allergy or asthma symptoms or breathing difficulties if inhaled. May cause an allergic skin reaction. Suspected of causing genetic defects. Suspected of causing cancer. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children. May cause damage to blood system, heart from repeated oral exposure at high doses.

**HANDLING PRECAUTIONS** Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. In case of inadequate ventilation wear respiratory protection. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves and protective clothing. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do NOT breathe mist/vapours/spray. Avoid contact during pregnancy/ while nursing. Avoid release to the environment. Very toxic to aquatic life. Harmful to the soil environment. Toxic to terrestrial vertebrates. Toxic to terrestrial invertebrates.

**FIRST AID** If medical advice is needed, have product container or label on hand. IF SWALLOWED: Call a POISON CENTRE or doctor/ physician if you feel unwell. Do NOT induce vomiting. IF INHALED: If breathing is difficult, remove to fresh air and keep at rest position comfortable for breathing. If experiencing respiratory symptoms: call a POISON CENTRE or doctor/ physician. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do so. Continue rinsing. Get medical advice/attention. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/ attention. Wash contaminated clothing before re-use. If exposed, or feel unwell, or concerned call a POISON CENTRE or get medical advice/ attention. **National Poisons Centre 24 Hour Phone : 0800 POISON (0800 764 766)**

**WARNINGS AND CONTRAINDICATIONS** • Please note that this product is in a concentrated form • Do not use in lambs under 20kg or under 4 weeks of age • Do not use in calves under 100kg • Do not use with milk or milk-feeders or while calves are suckling • Do not use on animals which are severely stressed or in ill health • Do not exceed stated dose or frequency without consulting veterinarian • Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity, so estimate live weights carefully. Dehydrated animals may be more susceptible to toxicity • Do not treat at intervals of less than 3 weeks, due to selenium content • Do not use at the same time as any other selenium fertilizer, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian • Fatal interactions may occur between levamisole and organophosphate dips • May be used in pregnant animals at recommended dose rates.



Cut along dotted line.



Registered to:  
Donaghys Ltd, 16 Sheffield Crescent,  
Harewood, Christchurch  
Freephone: 0800 942 006  
[www.donaghys.com](http://www.donaghys.com)

EPA Approval number HSR100758.  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions  
Registered pursuant to the ACVM Act 1997: A011650.  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.



UN3082  
ENVIRONMENTALLY HAZARDOUS SUBSTANCE LIQUID N.O.S.  
(CONTAINS Abamectin, Selenium)  
Packaging Group: III  
Hazchem: 3Z

Batch:  
Expiry:

ZPLAB945 MAR23 PNJ0316165-1 VERZ