

# Moxam<sup>®</sup>

## Sheep

### 🕒 SINGLE ACTING ORAL DRENCH FOR SHEEP

For the treatment of gastrointestinal and pulmonary nematodes and itch mite of sheep. Donaghys MOXAM Sheep contains moxidectin, a member of the milbemycin / avermectin family of anthelmintics. It is effective against sensitive strains of gastrointestinal roundworms, lungworm and itch mite.

NET CONTENTS  
10 Litres

**WARNING**  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

**ACTIVE CONSTITUENTS**  
Each mL contains 2mg of moxidectin.  
Also contains Benzyl Alcohol <10%  
w/v and BHT <1% w/v.

**DONAGHYS**  
PERFORMANCE FIRST

# Moxam<sup>®</sup>

## Sheep

### 🕒 SINGLE ACTING ORAL DRENCH FOR SHEEP

For the treatment of gastrointestinal and pulmonary nematodes and itch mite of sheep. Donaghys MOXAM Sheep contains moxidectin, a member of the milbemycin / avermectin family of anthelmintics. It is effective against sensitive strains of gastrointestinal roundworms, lungworm and itch mite.

NET CONTENTS  
10 Litres

**WARNING**  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

**ACTIVE CONSTITUENTS**  
Each mL contains 2mg of moxidectin.  
Also contains Benzyl Alcohol <10%  
w/v and BHT <1% w/v.

**DONAGHYS**  
PERFORMANCE FIRST

Philstic Labels Ltd accepts no legal responsibility for errors in text printed on our labels that falls under the recommended sizing guideline of 6pt.

425mm wide

**YOUR PROOF**

Job: 354408  
Name: 01 Moxam Sheep 10L Front  
Client: Donaghys Limited  
Date: 12-12-2022

CHECK A PANTONE BOOK FOR TRUE COLOUR REPRESENTATION

- OK to proceed  
 Further proof required

sign: \_\_\_\_\_ date: \_\_\_\_\_

No responsibility is taken for errors not noted on this proof  
**EXAMINE THIS ART VERY CAREFULLY AND RETURN  
THE SIGNED PROOF TO US FOR THE JOB TO PROCEED**

**PHILSTIC**<sup>®</sup> Philstic Labels Limited  
Email: prepress@philstic.co.nz  
Phone: +64 9 970 1430  
Fax: +64 9 630 0957

DIELINE

Yellow

Cyan

Magenta

Black

100mm deep

Donaghys MOXAM Sheep contains moxidectin, a member of the milbemycin /ivermectin family of anthelmintics. It is effective against sensitive strains of the following parasites:

**Gastrointestinal roundworms (adult and immature stages of):**

- Haemonchus contortus* (incl. inhibited or hypobiotic larvae)
- Ostertagia circumcincta* (incl. inhibited or hypobiotic larvae)
- Ostertagia trifurcata* (incl. inhibited or hypobiotic larvae)
- Trichostrongylus axei* (incl. inhibited or hypobiotic larvae)
- Trichostrongylus colubriformis* (incl. inhibited or hypobiotic larvae)
- Trichostrongylus vitrinus* (incl. inhibited or hypobiotic larvae)

- Nematodirus spathiger*
- Nematodirus filicollis*
- Cooperia curticei* (adult only)
- Cooperia oncophora* (adult only)
- Cooperia punctata* (adult only)
- Oesophagostomum venulosum* (adult only)
- Chabertia ovina*
- Trichuris ovis* (adult only)
- Strongyloides papillosus* (adult only)

**Lungworm:**  
Dictyocaulus filaria (adult and immature)

**External parasites:**  
Psorergates ovis (itch mite)  
Resistance may develop to any anthelmintic. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using parasite treatment.

**PERSISTENT ACTIVITY** Donaghys MOXAM Sheep when used at the recommended dose rate, prevents reinfection with *Haemonchus contortus* for 35 days and *Ostertagia circumcincta* for at least 21 days following a single oral dose.

**DIRECTIONS FOR USE** By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

FOR ORAL USE ONLY. SHAKE WELL BEFORE USE.

Use 1mL per 10kg live weight (equivalent to 0.2mg moxidectin per kg live weight). Administer using standard drenching equipment. Check accuracy of drench gun regularly. Correct drenching technique should be used. A representative sample of animals should be weighed before treatment. Draft animals into lines of similar body weight, then dose according to weight of the heaviest animal in the line. When there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing. Do not underdose.

**CONCURRENT TREATMENTS** Safe to use concurrently with other treatments in sheep including mineral supplements, fluke treatments and vaccines.

**WITHHOLDING PERIODS** It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**MILK:** Milk intended for human consumption must be discarded during treatment and for 35 days following the last treatment.  
**MEAT:** Animals producing meat and offal for human consumption must not be sold for slaughter either during or within 10 days of the cessation of the last treatment.

**STORAGE** Store below 25°C.

**DISPOSAL** Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Do not use container for any other purpose.

**WARNING**

**READ ENTIRE LABEL BEFORE USE**

May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child from repeated oral exposure. May cause harm to breast-fed children from repeated oral exposure.

**HANDLING PRECAUTIONS** Avoid skin contact. When handling wear protective gloves. Do not eat, drink or smoke when using this product. Do not breathe mist / vapours / spray. Avoid contact during pregnancy/while nursing. Wash thoroughly after handling. Contaminated work clothing should not be allowed out of workplace.

**FIRST AID** For advice contact the National Poisons Centre - 0800 POISON (0800 764 766) - or a doctor, immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

**ENVIRONMENTAL PROTECTION** Very toxic to aquatic organisms with long lasting effects. Harmful to the soil environment. Harmful to terrestrial invertebrates. Avoid contamination of any water supply with product or empty container. Avoid release to the environment.

Registered pursuant to the ACVM Act 1997, No A10378. See www.foodsafety.govt.nz for registration conditions.



DOSE RATE		
Bodyweight	Dose	Doses per 10L pack
16-20kg	2mL	5000
21-25kg	2.5mL	4000
26-30kg	3mL	3333
31-35kg	3.5mL	2857
36-40kg	4mL	2500
41-45kg	4.5mL	2222
46-50kg	5mL	2000
51-55kg	5.5mL	1818
56-60kg	6mL	1666
61-65kg	6.5mL	1538
66-70kg	7mL	1428
71-75kg	7.5mL	1333
76-80kg	8mL	1250

For animals over 80kg body weight use at a rate of 1mL per 10kg.



Registered to:  
Donaghys Ltd, 16 Sheffield Crescent,  
PO Box 20 449, Christchurch  
Freephone: 0800 942 006  
www.donaghys.com

**DONAGHY'S**  
PERFORMANCE FIRST

**YOUR PROOF**  
Job: 354408  
Name: 02 Moxam Sheep 10L Back  
Client: Donaghys Limited  
Date: 14-12-2022

CHECK A PANTONE BOOK FOR TRUE COLOUR REPRESENTATION

OK to proceed  
 Further proof required  
sign: \_\_\_\_\_ date: \_\_\_\_\_

No responsibility is taken for errors not noted on this proof  
**EXAMINE THIS ART VERY CAREFULLY AND RETURN THE SIGNED PROOF TO US FOR THE JOB TO PROCEED**

**PHILSTIC** Philstic Labels Limited  
Email: prepress@philstic.co.nz  
Phone: +64 9 970 1430  
Fax: +64 9 630 0957

- DIELINE
- Yellow
- Cyan
- Magenta
- Black

100mm deep

Philstic Labels Ltd accepts no legal responsibility for errors in text printed on our labels that falls under the recommended sizing guideline of 6pt.

425mm wide