

**DONAGHYS**

# TwinAL **Cattle**

**DOUBLE ACTIVE ORAL DRENCH FOR CATTLE**

For the control of roundworms, tapeworm, lungworm and adult liver fluke in cattle, including roundworms resistant to benzimidazole or levamisole.

NET CONTENTS  
**10 Litres**

**ACTIVE INGREDIENTS**  
100 g/L Albendazole and  
75 g/L Levamisole Hydrochloride.

**DONAGHYS**  
PERFORMANCE FIRST

**DANGER**  
KEEP OUT OF REACH OF CHILDREN  
READ LABEL BEFORE USE  
FOR ANIMAL TREATMENT ONLY

**DONAGHYS**

# TwinAL **Cattle**

**DOUBLE ACTIVE ORAL DRENCH FOR CATTLE**

For the control of roundworms, tapeworm, lungworm and adult liver fluke in cattle, including roundworms resistant to benzimidazole or levamisole.

NET CONTENTS  
**10 Litres**

**ACTIVE INGREDIENTS**  
100 g/L Albendazole and  
75 g/L Levamisole Hydrochloride.

**DONAGHYS**  
PERFORMANCE FIRST

**DANGER**  
KEEP OUT OF REACH OF CHILDREN  
READ LABEL BEFORE USE  
FOR ANIMAL TREATMENT ONLY

**READ ENTIRE LABEL BEFORE USE**

**INDICATIONS** For the control of sensitive mature and immature haemonchus, ostertagia, nematodirus, cooperia, strongyloides, bunostomum, oesophagostomum, chabertia that may be resistant to benzimidazole or levamisole. Also for the control of tapeworm, lungworm and adult liver fluke in cattle.

Resistance can develop to any anthelmintic.

Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

**HANDLING PRECAUTIONS** Harmful if swallowed. May cause an allergic skin reaction. Suspected of causing genetic defects. May damage fertility or the unborn child. May cause damage to organs. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves/clothing. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. Do not breathe dust/fume/gas/mist/ vapours/spray.

**FIRST AID** If medical advice is needed, have product container or label at hand.

**IF SWALLOWED:** Call a POISON CENTER or doctor/physician if you feel unwell. Rinse mouth.

**IF ON SKIN:** Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse.

If exposed or concerned, or feel unwell: Get medical advice/attention.

For advice, contact the National Poisons Centre – 0800 POISON (0800 764 766) or a doctor immediately.

**ENVIRONMENTAL PROTECTION** Harmful to terrestrial vertebrates. Avoid release to the environment and contamination of any water supply with product or empty container. Collect spillage.

**PRECAUTIONS** Doses three or more times those recommended can cause symptoms of Levamisole toxicity, so estimate liveweights carefully. Animals under stress due to advanced pregnancy, adverse weather conditions, poor nutrition, prolonged yarding or dehydration may be more susceptible to toxicity.

Fatal interactions may occur between Levamisole and organophosphate dips.

Do not exceed the stated dose volume.

**DOSAGE** Cattle 1 mL per 10 kg bodyweight.

Bodyweight	Dose	Dose per 10 L Pack
60 kg	6 mL	1666
61-70 kg	7 mL	1428
71-80 kg	8 mL	1250
81-90 kg	9 mL	1111
91-100 kg	10 mL	1000
101-110 kg	11 mL	909
111-120 kg	12 mL	833
121-130 kg	13 mL	769
131-140 kg	14 mL	714
141-150 kg	15 mL	666
151-160 kg	16 mL	625
161-170 kg	17 mL	588
171-180 kg	18 mL	555

Bodyweight	Dose	Dose per 10 L Pack
181-190 kg	19 mL	526
191-200 kg	20 mL	500
201-210 kg	21 mL	476
211-220 kg	22 mL	454
221-230 kg	23 mL	434
231-240 kg	24 mL	416
241-250 kg	25 mL	400
251-260 kg	26 mL	384
261-270 kg	27 mL	370
271-280 kg	28 mL	357
281-290 kg	29 mL	344
291-300 kg	30 mL	333

For animals heavier than 300 kg add 1 mL per 10 kg bodyweight.

**DIRECTIONS FOR USE** By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than directed on the label.

Draft animals into lines of similar bodyweight. A representative sample of animals should be weighed before treatment. Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing. Correct drenching techniques should be used. Check accuracy of drench guns regularly.

**DISPOSAL** Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill facility. Triple rinse container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at approved landfill facility. Do not use container for any other purpose.

**STORAGE** Store locked up below 25°C. Away from sunlight. Do not freeze.

**WITHHOLDING PERIODS** It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**MEAT:** Animals producing meat or offal for human consumption must not be sold for slaughter during or within 14 days of cessation of last treatment.

**MILK:** Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID N.O.S. (Contains: Albendazole), UN3082

Registered pursuant to the ACVM Act 1997, No A11778. See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, HSR100758. See www.epa.govt.nz for controls.

Registered to:  
Donaghys Ltd, 16 Sheffield Crescent, Harewood, Christchurch  
Freephone: 0800 942 006 | www.donaghys.com

**DONAGHYS**  
PERFORMANCE FIRST



Batch:  
Expiry: